## Food and Drug Administration, HHS

A article), 50 to 720 milligrams of lasalocid, and 45 to 180 milligrams of tylosin per pound to a ration of non-medicated feed.

- (7) Amount. 0.5 milligram per head per day.
- (i) *Indications for use*. For suppression of estrus (heat).
- (ii) Limitation. Heifers intended for breeding. Do not exceed 24 days of feeding. Administer 0.5 to 2.0 pounds per head per day of Type C feed containing 0.25 to 1.0 milligram of melengestrol acetate per pound to provide 0.5 milligram of melengestrol acetate per head per day. Melengestrol acetate as provided by No. 000009 in §510.600(c) of this chapter.
- (8) Amount. Melengestrol acetate, 0.25 to 0.5 milligram per head per day, plus oxytetracycline, 75 milligrams per head per day.
- (i) Indications for use. For increased rate of weight gain, improved feed efficiency, suppression of estrus (heat), and reduction of liver condemnation due to liver abscesses.
- (ii) Limitations. Heifers fed in confinement for slaughter. Add at the rate of 0.5 to 2.0 pounds per head per day a medicated feed (liquid or dry) containing 0.125 to 1.0 milligram of melengestrol acetate per pound to a feed containing 6 to 10 grams of oxytetracycline per ton; or add at the rate of 0.5 to 2.0 pounds per head per day a dry medicated feed containing 0.125 to 1.0 milligram of melengestrol acetate plus 37.5 to 150 milligrams of oxytetracycline per pound to provide 0.25 to 0.5 milligram of melengestrol acetate and 75 milligrams of oxytetracycline per head per day. Liquid melengestrol acetate may not be mixed with oxytetracycline in a common liquid feed supplement. Melengestrol acetate as provided by 000009, oxytetracycline by 000069, in §510.600(c) of this chapter.

[42 FR 28535, June 3, 1977, as amended at 46 FR 18967, Mar. 27, 1981; 49 FR 39842, Oct. 11, 1984; 50 FR 1843, Jan. 14, 1985; 50 FR 16228, Apr. 25, 1985; 51 FR 7398, Mar. 3, 1986; 51 FR 36222, Oct. 9, 1986; 52 FR 2685, Jan. 26, 1987; 52 FR 33804, Sept. 8, 1987; 53 FR 27959, July 26, 1988; 55 FR 31828, Aug. 6, 1990; 57 FR 5053, Feb. 12, 1992; 59 FR 41241, Aug. 11, 1994; 60 FR 11028, Mar. 1, 1995; 62 FR 8373, Feb. 25, 1997; 62 FR 14304, Mar. 26, 1997; 63 FR 27845, May 21, 1998; 63 FR 41191, Aug. 3, 1998; 63 FR 66018, Dec. 1,

## §558.348 Mibolerone.

- (a) Approvals. To No. 000009 in \$510.600(c) of this chapter for a canned dog food, each  $6\frac{1}{2}$  ounce can containing 30 or 60 micrograms of mibolerone.
- (b) Conditions of use—(1) Amount. 30 micrograms for animals weighing up to 25 pounds; 60 micrograms for animals weighing 26 to 50 pounds; 120 micrograms for animals weighing 51 to 100 pounds; 180 micrograms for animals weighing over 100 pounds, or German Shepherds or German Shepherd mix weighing 30 to 80 pounds.
- (2) *Indications for use*. For the prevention of estrus (heat) in adult female dogs not intended primarily for breeding purposes.
- (3) Limitations. Administer daily at least 30 days before expected initiation of heat and continue as long as desired, but for not more than 12 months. Mibolerone should not be used in bitches before first estrous period or in purebred Bedlington terriers. It is not intended for animals being used primarily for breeding purposes. Use orally in adult female dogs only. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[47 FR 6617, Feb. 16, 1982]

## § 558.355 Monensin.

- (a) Specifications. Monensin, as the base or the sodium salt, contains a minimum of 90 percent monensin activity derived from monensin A and a minimum of 95 percent derived from monensin A plus B. Using thin layer chromatography, the  $R_f$  value must be comparable to a reference standard (the  $R_f$  value is the distance the spots travel from the starting line divided by the distance the solvent front travels from the starting line), or, using High Performance Liquid Chromatography, the factor distribution of monensin Factor A or B is calculated as the percentage of total biopotency of all peaks. The loss on drying is not more than 10 percent when dried in vacuum at 60 °C for 2 hours.
- (b) Approvals. Approvals for Type A medicated articles containing the specified levels of monensin activity granted to firms identified by sponsor numbers in §510.600(c) of this chapter for